

# Basic quantum field theory

By J.A.J. van Leunen

Last modified: 7 oktober 2017

## Abstract

In the eighteenth century, scientists discovered the ingredients of basic quantum field theory. In those times quantum physics played no role. In the twentieth century, these ingredients were forgotten and stayed ignored.

This paper introduces two categories of super-tiny dark objects that represent the most basic field quanta. Warps represent a tiny bit of energy. Clamps represent a tiny bit of mass. Observers cannot perceive these objects as individual items. The objects are the tiny dark objects that science is still missing. The LHC and its successors will never be able to detect them.

The paper shows that one-shot triggers can expand continuums and the expansion can represent an amount of mass.

## Introduction

Quantum field theory requires a continuum that can be deformed or vibrated and actuators that cause this deformation or vibration. Next, the activity of these actuators must be quantized. Thus, the strength of the deformation or vibration occurs in a set of fixed values. Deformation or vibration can be temporarily or persistent.

## Field dynamics

A function for which both the parameter space and the target space are multi-dimensional describes a continuum. Dynamics requests a progression parameter, and the spatial part requests a multi-dimensional spatial parameter. Quaternions have the advantage that they combine storage for the progression part, and the spatial part and quaternionic calculus defines a multiplication procedure for the combination of the two. Quaternions can also store the scalar part and the vector part of the target value of a quaternionic function. Quaternions can describe the behavior of dynamic fields via [quaternionic differential calculus](#). Partial second order differential equations describe the interaction between point-like artifacts and quaternionic continuums.

The combination of a quaternionic infinite dimensional separable Hilbert space and its unique non-separable companion Hilbert space that embeds its separable partner offers the playground where this interaction can take place. This playground stores separate quaternions in eigenspaces of operators that reside in the separable Hilbert space and can store quaternionic continuums as eigenspaces of operators that reside in the non-separable Hilbert space. Quaternionic functions define these continuums.

A subspace that scans this base model as a function of a selected progression value represents the static status quo of the model and splits it between a historical part, the current static status quo, and a future part.

The embedding maps the discrete quaternions onto an embedding continuum. The embedding process occurs inside the scanning subspace.

## Solutions of the differential equations

The dynamic solutions of the homogeneous second order partial differential equations do not occur spontaneously. Actuators that determine what kind of solution results generate them. For example, a periodic harmonic actuator causes wave solutions of a homogeneous second order partial differential equation, which is therefore known as the wave equation. Also, another, quite similar homogeneous second order partial differential equation exist that does not offer waves as its solutions. This equation splits into two first order partial differential equations. Both homogeneous second order partial differential equations offer solutions in the form of shock fronts that one-shot actuators trigger. These solutions occur in two versions. **Warps** are one-dimensional shock fronts that during travel keep their amplitude. **Clamps** are spherical shock fronts that quickly fade away because their amplitude diminishes as  $1/r$  with distance  $r$  from the trigger location. In the meantime, clamps integrate into the Green's function of the carrier field, which means that they temporarily deform the carrier. Warps carry a standard bit of energy and clamps carry a standard bit of mass. These facts make them the most **basic quanta** of the carrier field. Point-like actuators trigger the warps and the clamps. That does not mean that more powerful triggers cannot trigger correspondingly larger shock fronts. An isotropic explosion may cause a gravitational spherical shock front. Every triggered vibration may cause a more permanent expansion of the embedding continuum and it may represent an amount of mass.

## Super-tiny dark objects

Warps and clamps form two categories of super-tiny objects that in separation cannot be perceived. Only organized in huge collections, these objects become observable. For example, if emitted at equidistant instants, the warp strings become a frequency, and if these strings obey the Einstein-Planck relation, then the strings implement the functionality of photons.

If recurrently regenerated by dense and coherent swarms of hop landing location triggers, the clamps become noticeable as elementary particles.

Less coherent assemblies of warps can create a noticeable amount of **dark energy**. Less coherent assemblies of clamps can create a noticeable amount of **dark mass**.

Elementary particles are elementary modules. Together these elementary modules generate all other modules and the modules construct modular systems.

## Ensuring coherence

A private mechanism that applies a stochastic process, which owns a characteristic function generates a hopping path and a hop landing location swarm. The characteristic function acts as a displacement generator and ensures that the process generates a coherent swarm, which moves as a single unit. The location density distribution of the swarm is the Fourier transform of the characteristic function and equals the squared modulus of the wavefunction of the object that the swarm represents.

The generated swarms represent elementary modules. They show both particle and wave behavior. The characteristic function of the stochastic process explains the wave behavior.

Elementary modules reside on private platforms that own a private parameter space that a version of the quaternionic number system generates. The platforms float over a background parameter space, which the version of the quaternionic number system that the Hilbert spaces use to define their inner product generates. The differences in ordering symmetry between parameter spaces give

rise to symmetry-related charges. These charges locate at the geometrical centers of the platforms and produce symmetry-related fields.

The elementary modules inherit the properties of the platforms on which they reside. In this way, a range of different elementary module types exist.

With every new embedded location the stochastic mechanisms expand the embedding continuum.

## Modules and spectral binding

Together the elementary modules constitute all other modules and the modules constitute modular systems.

Also here, stochastic processes that own a characteristic function generate the footprints of modules. Therefore, the module also moves as a single unit.

The characteristic function of the module equals the superposition of the characteristic functions of the components of the module. The superposition coefficients determine the internal locations of the components. These coefficients may oscillate.

The superposition installs a very strong kind of spectral binding.

Gravity and attractive symmetry-related charges may add to the effect of spectral binding.

## Recurrent regeneration

Clamps quickly fade away and therefore, clamps require the recurrent regeneration of the objects that they generate. The regeneration cycle determines how much clamps, thus how much mass the generated object owns. Thus, the current static status quo is surrounded by a particle regeneration fabric.

## History

The solutions of the wave equation are known for more than two and a half centuries [1]. In those times physicists were not aware of the quantization of space, but some awareness was growing about the quantization of wave packages. The shock fronts are not waves. They do not feature a frequency. Wave packages disperse when they move. Shock fronts do not disperse.

It is strange that during the development of quantum physics the shock fronts escaped the attention of the early quantum physicists. Otherwise, basic quantum field theory would have become a straightforward part of quantum theory.

Currently LIGO and Virgo are observing gravitational waves. The cause is the end of the mutual rotation around a common center of two black holes. The subsequent explosion need not be isotropic. Isotropic one-shot triggers cause spherical shock fronts. Still the phenomenon can deform our living space and may permanently expand this continuum. This phenomenon may represent an amount of mass because it permanently deforms the carrier continuum.

## Mathematics

Partial quaternionic differential equations that apply the quaternionic nabla  $\nabla$  describe the interaction between a field and a point-like artifact [2].

$$\nabla \equiv \{\partial/\partial\tau, \partial/\partial x, \partial/\partial y, \partial/\partial z\}$$

$$\nabla \equiv \{\partial/\partial x, \partial/\partial y, \partial/\partial z\}$$

$$\nabla_r \equiv \partial/\partial\tau$$

$\tau$  is progression or proper time.

In the quaternionic differential calculus, differentiation with the quaternionic nabla is a quaternionic multiplication operation:

$$c = c_r + \mathbf{c} = ab \equiv (a_r + \mathbf{a}) (b_r + \mathbf{b}) = a_r b_r - \langle \mathbf{a}, \mathbf{b} \rangle + \mathbf{a} b_r + a_r \mathbf{b} \pm \mathbf{a} \times \mathbf{b}$$

Here the real part gets subscript  $r$  and the imaginary part is written in bold face.

The right side covers five different terms.

$\langle \mathbf{a}, \mathbf{b} \rangle$  is the inner product.

$\mathbf{a} \times \mathbf{b}$  is the external product.

$\pm$  indicates the choice between right and left handedness.

Now the partial differential equation that describes the first order behavior of a continuum is given by:

$$\Phi = \phi_r + \mathbf{\Phi} = \nabla\psi \equiv (\nabla_r + \nabla) (\psi_r + \boldsymbol{\psi}) = \nabla_r \psi_r - \langle \nabla, \boldsymbol{\psi} \rangle + \nabla \psi_r + \nabla_r \boldsymbol{\psi} \pm \nabla \times \boldsymbol{\psi}$$

$$\phi_r = \nabla_r \psi_r - \langle \nabla, \boldsymbol{\psi} \rangle$$

$$\mathbf{\Phi} = \nabla \psi_r + \nabla_r \boldsymbol{\psi} \pm \nabla \times \boldsymbol{\psi}$$

$\langle \nabla, \boldsymbol{\psi} \rangle$  is the divergence of  $\boldsymbol{\psi}$

$\nabla \psi_r$  is the gradient of  $\psi_r$

$\nabla \times \boldsymbol{\psi}$  is the curl of  $\boldsymbol{\psi}$

$$\mathbf{E} = -\nabla \psi_r - \nabla_r \boldsymbol{\psi}$$

$$\mathbf{B} = \nabla \times \boldsymbol{\psi}$$

Double differentiation leads to the second order partial differential equation:

$$\rho = \nabla^* \Phi = (\nabla_r - \nabla) (\nabla_r + \nabla) (\psi_r + \boldsymbol{\psi}) = (\nabla_r \nabla_r + \langle \nabla, \nabla \rangle) (\psi_r + \boldsymbol{\psi}) = \rho_r + \mathbf{J}$$

This equation splits into two first order partial differential equations  $\Phi = \nabla\psi$  and  $\rho = \nabla^*\Phi$ .

$$\rho_r = \langle \nabla, \mathbf{E} \rangle$$

$$\mathbf{J} = \nabla \times \mathbf{B} - \nabla_r \mathbf{E}$$

$$\nabla_r \mathbf{B} = -\nabla \times \mathbf{E}$$

Two quite similar second order partial differential operators exist. The first is described above.

$$(\nabla_r \nabla_r + \langle \nabla, \nabla \rangle) \psi = \rho$$

This is still a nameless equation.

The second is the quaternionic equivalent of d'Alembert's operator ( $\nabla_r \nabla_r - \langle \nabla, \nabla \rangle$ ). It defines the quaternionic equivalent of the well-known wave equation.

$$(\nabla_r \nabla_r - \langle \nabla, \nabla \rangle) \psi = \varphi$$

Both second order partial differential operators are Hermitian differential operators.

## Solutions

### Waves

$$f(\tau, \mathbf{x}) = a \exp(i \omega(c\tau - |\mathbf{x} - \mathbf{x}'|)); \quad c = \pm 1$$

solves  $\nabla_r \nabla_r f = \langle \nabla, \nabla \rangle f = -\omega^2 f$

### Warps

$$\psi = g(\mathbf{x} \pm \tau)$$

### Clamps

$$\psi = g(r \pm \tau)/r$$

Except for  $\phi_r = \nabla_r \psi_r - \langle \nabla, \psi \rangle$  the above equations remind of Maxwell equations. In common quantum field theory, the exception is treated as a gauge. Maxwell equations use coordinate time, while the quaternionic differential equations apply proper time.

## References

[1] [https://en.wikipedia.org/wiki/Wave\\_equation#General\\_solution](https://en.wikipedia.org/wiki/Wave_equation#General_solution)

[2]

[https://en.wikiversity.org/wiki/Hilbert\\_Book\\_Model\\_Project/Quaternionic\\_Field\\_Equations](https://en.wikiversity.org/wiki/Hilbert_Book_Model_Project/Quaternionic_Field_Equations)